

Arman 2

by Cek Turnitin

Submission date: 06-Apr-2022 07:05AM (UTC+0000)

Submission ID: 1803195647

File name: Inter,_Copernicus_Arman_Jendri.pdf (366.7K)

Word count: 4125

Character count: 25139

Endophytic Bacteria and Fungi from Indonesian Medicinal Plants with Antibacterial, Pathogenic Antifungal and Extracellular Enzymes Activities: A Review

Jendri Mamangkey^{1*}, Lucas William Mendes², Arman Harahap³, Dana Briggs⁴, Cengiz Kayacilar⁵

9

¹ Department of Biology Education, Faculty of Education and Teacher Training, Universitas Kristen Indonesia, Jakarta Timur 13630, Jakarta, Indonesia.

13

² Center for Nuclear Energy in Agriculture, University of Sao Paulo, Piracicaba-SP, Brazil.

³ Department of Biology Education, Faculty of Education and Teacher Training, Universitas Labuhanbatu, Indonesia.

⁴ California State University, Monterey Bay, Marina, United States.

⁵ Niğde Ömer Halisdemir University, Turkey.

*Corresponding Author:

Email: jendri.mamangkey@uki.ac.id

Abstract.

The habitat of endophytic bacteria and fungi in plant tissues makes it automatically used by plants to protect themselves against pathogenic microorganisms. Endophytic bacteria and fungi have been applied in various fields, including agriculture, pharmacy, medicine, and biotechnology. This review was carried out to summarize recent studies focusing on the diversity of endophytic bacteria derived from traditional medicinal plants that have antibacterial activity and extracellular enzymes. This review article was prepared and written by referring to literature studies collected from online journal publications of endophytic bacteria focused on *Vernonia anthelmintic*, *Saurauia scaberrinae*, and *Ki rinyuh* (*Chromolaena odorata*) plants. The results revealed that endophytic bacteria obtained from *Vernonia anthelmintic* plant belong to the bacterial genera *Micrococcus*, *Bacillus*, *Pseudomonas*, *Stenotrophomonas*, and *Pantoea* which possess the same biological properties as the host plant, producing antibacterial activity. The *Saurauia scaberrinae* plant successfully isolated endophyte fungus *Phoma* sp. which produces phomodione (4) and cercosporamide. Phomodione (4) exhibit inhibitory activity against *S. aureus*, *P. ultimum*, *S. sclerotiorum* and *R. solani*. Cercosporamide exhibits inhibitory activity against *S. aureus*. Endophytic bacteria were isolated from *Ki rinyuh*, including BECB3, BECB 4, BECA 8, BECA 5, BECA 1, and BECA 10 isolates which have extracellular enzyme activities such as β -amylase, α -amylase, cellulase, chitinase, and protease. Understanding endophytic bacteria in medicinal plants can help researchers them effectively as a biotechnological product.

Keywords: Endophytic bacteria, medicinal plants, antibacterial, extracellular enzymes

I. INTRODUCTION

Research on endophytic bacteria and fungi derived from medicinal plant organs in past decades is continuously indicating an escalation graph. Such increase was affected by the scientists' movement to find bioactive compounds of endophytic bacteria and fungi that are similar to their host plant compounds. Furthermore, the term 'endophyte' should be noted as the name proposed by Heinrich Anton de Bary in 1866 which was taken from the Greek words *endon* (inside) and *phyte* (plant) [1].

Endophytes are generally related to nonpathogenic bacteria and fungi inhabiting plant tissues, but are sometimes also used to refer to mycorrhizae found in plant roots. The positive effects due to the presence of endophytes for the plants include increasing the health, size, and continuity of plant growth. Endophytic bacteria and fungi are currently categorized as potential sources of new natural products for their exploitation in the field of medicine, agriculture, and industry. Endophytic bacteria isolated from traditional medicinal plants have been reported to produce antibacterial, antifungal, and antiseptic compounds [2,3]. Furthermore, the secondary metabolites from endophytic fungal cultures have been previously proven to have cytotoxic, antimicrobial, antiviral, and anticancer activities [4,5,6,7,8,9,10,11,12]. In addition, plants that have significant biological activity have the potential to hide endophytes with the same bioactivity [11,13]. In this case, for example, endophytic bacteria isolated from medicinal plants of *Solanum distichum*, *Matricaria chamomilla*, *Calendula officinalis* [14], and *Hypericum perforatum* [15] show excellent antibacterial and antifungal activity.

In addition, endophytic fungi isolated from leaves of the endemic plant of *Zygodium Mandeville*, which is *Cladosporium Cladosporioides*, has the best antimicrobial activity causing an inhibition ranging from 20.7 to 25.7 mm against phytopathogenic bacteria and fungi [16]. Endophytes have an antimicrobial activity that can produce metabolites, such as *celastramycins* [17], *kakadumycins* [18], and *javanicin* [19]. Reports from other previous research further revealed that endophytic bacteria and fungi have extracellular enzyme activity [20,21]. Endophytic bacteria and fungi synthesize extracellular enzymes as a resistance mechanism against disease, which helps to fend off pathogens and obtain food from their host plants. The extracellular enzymes that are commonly produced by endophytic bacteria and fungi are pectinase, amylase, cellulase, lipase, and protease. *Pseudomonas aeruginosa* was successfully isolated from the leaves of *Anredera cordifolia*, producing proteases, amylase, esterase, and cellulase [22]. *Bacillus subtilis* P4 isolated from the *Pseudobrickellia brasiliensis* plant has protease and esterase activity [23]. Furthermore, the endophytic bacterium *Lysinibacillus xylanilyticus* has also been isolated from the *Chiliadenus Montanus* plant and potentially produces cellulase [24]. *Vernonia anthelmintica*, *Saurauia scaberrinae*, and *Chromolaena odorata* are categorized as medicinal plants. The *Vernonia* genus, which belongs to the *Asteraceae* family with 1000 species, has been widely used for treating diabetes, parasitic worms, asthma, inflammation, wounds, skin diseases, malaria, cancer, and digestive disorders [25,26].

Various biological effects of *V. anthelmintica* plant extracts have also been reported, including anti-inflammatory, antimicrobial [27], antidiabetic and antihyperlipidemic activities [28]. Plants of *Saurauia* genus are widely distributed in Indonesia. One species that is considered quite widely known is *Saurauia bracteosa* which grows widely in the area of Lake Toba, North Sumatra, Indonesia [29]. Furthermore, Silalahi [30] reported that the leaves of *Saurauia vulcanii* were used for

treating diarrhea, indigestion, and ethnomedicine for the Simalungun Batak sub-ethnic, in North Sumatra Province, Indonesia. Another species, *Saurauia laevigata*, is a 'honduran' medicinal plant used to treat stomachache, in addition to *Saurauia scaberrinae*, which is found in Papua New Guinea, whose extracts *scaberrinae* contain alkaloid compounds [31]. Kirinyuh (*Chromolaena odorata*) plant is a wild plant easily found in our environment. The leaves of this plant contain several important compounds such as tannins, phenols, flavonoids, saponins, alkaloids, and steroids, while their essential oil contains α -pinene, cadinene, camphora, limonene, β -caryophyllene, and cadinol isomers [32]. Although the medicinal potential of these plants is well explored, unfortunately, research on endophytic bacteria and fungi from these plants has not been carried out optimally. Hence, the current review aims to explain the endophytic bacteria and fungi that have been studied from *Vernonia anthelmintica*, *Saurauia scaberrinae*, and *Chromolaena odorata* with the purpose of giving opportunities to the other researchers to conduct in-depth study on the plants that have not been studied yet. A better understanding of the relationship between endophytic microorganisms and their host medicinal plants will provide knowledge that can be exploited for the biotechnological production of better drugs from these plants.

II. METHODS

The writing and preparation of this article were carried out based on a literature review obtained from online textbooks, research results, and scientific journals that have been published online. The research data used in this study are categorized as secondary data. Keywords used in online searches include endophytic bacteria, endophytic fungi, *Vernonia anthelmintic*, *Saurauia scaberrinae*, and *Chromolaena odorata*. Data that have been collected were further cohered to describe endophytic bacteria and fungi in medicinal plants of *Vernonia anthelmintica*, *Saurauia scaberrinae*, and *Chromolaena odorata*.

III. RESULT AND DISCUSSION

Vernonia anthelmintica plant is categorized as *Asteraceae* family and is widely distributed in subtropical and tropical areas throughout Asia and Africa [33]. This plant growth can reach 90 cm and grows widely throughout India, mostly in places near landfills to a height of 1500 m [34,35]. This plant has been generally used as a traditional medicine to cure quite many diseases including asthma, wounds, inflammation, skin diseases, kidney problems, eye itching, and hiccups [35,36].

This medicinal plant has also been reported to have various pharmacological activities such as antimicrobial, anticancer, antidiabetic, anti-inflammatory, analgesic, antipyretic, as well as diuretic and larvicidal activities [37]. This plant is commonly used in rural and urban areas of Bangladesh for treating diabetes [38]. Through phytochemical studies, it has been discovered that *V. anthelmintica* contains fatty acids, steroids, flavonoids, sesquiterpene lactones, carbohydrates, and terpenes [33,37].



Fig 1. *Vernonia anthelmintica* plant. Rustamova et al. (2020)

Considering its benefits in the medical field, more research was conducted on *V. anthelmintica* plant by tracing its endophytic bacteria and fungi. In this c³² *V. anthelmintica* was collected from the Xinjiang Autonomous Forest Region of the People's Republic of China in previous research [39]. Thus, Rustamova et al. [39] successfully isolated 25 bacterial stirpes from *V. anthelmintica* plant tissue, the isolates were identified as *M. endophyticus* VERA1, *B. megaterium* VERA2, *P. chlororaphis* VERA3, *P. kilonensis* VERA4, *S. pavanii* VERA5, *B. endophyticus* VERA6, *S. maltophilia* VERA7, *P. ananatis* VERA8, *B. atrophaeus* VERA5. *S. pavanii* VERA5 and *P. kilenensis* VERA4, showing antimicrobial activity against *E. coli*. The eight bacterial isolates except *S. pavanii* VERA5 and *B. atrophaeus* VERA9 showed moderate or strong inhibition against *S. aureus*. Previously, Emiliani et al. [40] observed the endophytic bacterium *Stenotrophomonas maltophilia* from the medicinal plant of *Lavandula angustifolia* Mill. Janardhan and Vijayan [41] also observed *Pantoea ananatis* from the medicinal plant of *Lantana camara* from Malaysia. Based on these studies, endophytic microorganisms associated with medicinal plants can synthesize biologically active compounds and show the same biological activity as ³⁸ir host cells (medicinal plant tissues) [42]. Furthermore, Akinsanya et al. [43] stated that endophytic bacteria associated with medicinal plants produce bioactive compounds with antimicrobial activity similar to their hosts.

Endophytes can also benefit their hosts by producing various natural products known as secondary metabolites. In this case, the endophytes can improve plant performance and protect it against insect pests and diseases infected from the soil or induce abiotic stress tolerance in plants [44,45]. Therefore, endophytes have a great potential to be used as a safe and cost-effective alternative to biopesticides and can be

used as fertilizers that are considered constituents of sustainable agriculture. Furthermore¹⁹ the medicinal plant *Saurauia scaberrinae* was also successfully collected from the area of Papua New Guinea. Hoffman et al.¹⁹ [6] successfully found a furandione compound named phomodione (4), which³⁵ is produced by the endophytic fungus of *Phoma sp.* Phomodione (4) showed activity against *S. aureus* (MIC 1.6 µg/mL), *P. ultimum* (4–5 µg/mL), *S. sclerotiorum* (3–5 µg/mL), and *R. solani* (5– 8 µg/mL). As a reference, comparison was conducted between phomodione (4) activity and cercosporamide activity since their structure is similar. Cercosporamide is another natural product synthesized by *Phoma sp.* Cercosporamide, showing activity against *S. aureus* (MIC²⁷ g/mL). Not only that, cercosporamide can inhibit *Pythium ultimum* (3– 4 µg/mL), plant pathogenic fungi from oomycetes class which are responsible for various diseases in many plants. Cercosporamide can also inhibit *Sclerotinia sclerotiorum* (5–8 µg/mL) which belongs to the plant pathogenic fungi of the ascomycetes class and *Rhizoctonia solani* (8–10 g/mL) of the plant pathogenic fungi of the basidiomycetes class.



Fig 2. Leaves of *Saurauia vulcani* plant. Pasaribu et al. [29]

10

The endophytic fungus *Phoma sp.* was also successfully isolated from the medicinal plant *Artemisia princeps* Pamp. Lim et al. [47] reported two new derivatives of isochromanone compounds, (3S,4S)-3,8-dihydroxy-6-methoxy-3,4,5-trimethylisochroman-1-one and methyl (S)-18-hydroxy-6-methoxy-5-methyl-4a-(3-oxobutan-2-yl) benzoate isolated from endophytic fungus *Phoma sp.* PF2 culture. Similarly, da Silva et al. [48] successfully isolated *Phoma sp.* URM 7221 endophytic fungi from the leaves of *Schinus terebinthifolius*, *Phoma sp.* URM 7221 has antimicrobial activity against human pathogenic bacteria, which are *S. aureus* and Methicillin-Resistant *Staphylococcus aureus* (MRSA). *Phoma sp.* is predominantly found in the stems of *Artemisia annua*, contributing 50% of the total fungi isolated from the stems of *Artemisia annua* which have cellulase activity. Based on the results

of research conducted by Liu et al. [49] the cellulase activity of *Fusarium sp.* and *Phoma sp.* is significantly higher than *Humicola sp.* and *Alternaria sp.*

The endophytic bacteria of the medicinal plant *Chromolaena odorata* have also been previously studied. Mamangkey et al. [20] successfully reported 19 endophytic bacteria isolates from the roots, stems, and leaves of *Chromolaena odorata*. BECB3 isolates from stems and BECA8 were positive for their ability to produce β -amylase, α -amylase, cellulase, chitinase, and proteases. Most of the isolates produced proteases, except for isolates BECD3, BECD4, and BECA4. BECB3 produced significant hydrolytic enzyme activity, β -amylase (+++), α -amylase (++), cellulase (+++), chitinase (++) and protease (+++) compared to other isolates. In addition to bacteria, endophytic fungi were also successfully isolated from the leaves of *Chromolaena odorata*, which is CO-2AL3 isolate. The results of antimicrobial testing of the CO-2AL3 isolate extract showed antibacterial activity against *S. aureus*, *B. subtilis*, and *S. typhi* [50].



Fig 3. *Chromolaena odorata*

Gultom et al. [51] claimed that there were differences in the secondary metabolite compounds between young leaves and old leaves of kirinyuh (*Chromolaena odorata*) through gas chromatography. They found that there were more compounds obtained in old leaves of 22 bioactive compounds than the bioactive compounds found in young leaves, which are only 13 bioactive compounds. In this case, γ -sitosterol compounds are identified only in old leaves, while diisooctyl phthalate and Bis(2-ethylhexyl) phthalate, which belong to the saponin class are only found in young leaves. These bioactive compounds include flavonoids, terpenoids, saponins, phenols, alkaloids, and steroids [51].

IV. CONCLUSION

Endophytic bacteria and fungi co-evolved with plants and established a special relationship with their host, living inside the healthy tissues and being an important component of plant micro-ecosystems. In this review, we summarized recent studies on the diversity of endophytic bacteria and fungi derived from traditional medicinal plants. Briefly, endophytic bacteria from *V. anthelmintica* were identified as *M. endophyticus* VERA1, *B. megaterium* VERA2, *P. chlororaphis* VERA3, *P. kilonensis* VERA4, *S. pavanii* VERA5, *B. endophyticus* VERA6, *S. maltophilia* VERA7, *P. ananatis* VERA8, *B. atrophaeus* VERA9. *S. pavanii* VERA5 and *P. kilonensis* VERA4, which have antimicrobial activity against *E. coli*. Among these eight bacterial isolates, except *S. pavanii* VERA5 and *B. atrophaeus* VERA9, showed moderate or strong inhibition against *S. aureus*. Furthermore, endophytic fungi *Phoma sp.*

Was successfully isolated from the *Saurauia scaberrinae* plant, which produced *phomodione* (4) and showed activity against *S. aureus*, *P. ultimum*, *S. sclerotiorum* and *R. solani*. Cercosporamide is another natural product synthesized by *Phoma sp.* Cercosporamide which further showed activity against *S. aureus*. In this case, there were 19 isolates of endophytic bacteria derived from the roots, stems, and leaves of *Chromolaena odorata*. BECB3 isolates from stems and BECA8 were positive for their ability to produce β -amylase, α -amylase, cellulase, chitinase, and proteases. Endophytic fungi were also isolated from the leaves of *Chromolaena odorata*, which is CO-2AL3 isolate. In this case, antimicrobial activity was shown by the CO-2AL3 isolate extract against *S. aureus*, *B. subtilis*, and *S. typhi*. This report confirms that the bioactive compounds in medicinal plants are a habitat for specific endophytic bacteria and fungi that survive to live in medicinal plant tissues. However, all types of endophytic bacteria and fungi can live in these tissues, resulting in secondary metabolites produced by endophytic bacteria and fungi that are similar to the plants they live in.

REFERENCES

- [1] Bacon, C.W., White, J.F. (2000). *Microbial Endophytes*. New York, NY: Marcel Dekker.
- [2] Castillo, U.F., Strobel, G.A., Ford, E.J.F. (2002). Munumbicins, wide-spectrum antibiotics produced by *Streptomyces* NRRL 30562, endophytic on *Kennedia nigriscans*. *Microbiology* 148 (Pt 9): 2675-2685.
- [3] Berg, G., Hallmann, J. (2006). *Control of plant pathogenic fungi with bacterial endophytes*. In: Schulz B, Boyle C, Sieber TN, eds. *Soil Biology 9: Microbial root endophytes*. Berlin, Springer. pp 63 – 67.
- [4] Strobel, G., Daisy, B. (2003). Bioprospecting formicrobial endophytes and their natural products, *Microbiol. Mol. Biol. Rev.* 67: 491-502.

- [5] Strobel, G., Daisy, B., Castillo, U., Harper, J. (2004). Natural products from endophytic microorganisms, *J. Nat. Prod.* 67: 257-268.
- [6] Deshmukh, S.K., Gupta, M.K., Prakash, V., Saxena, S. (2018). Endophytic fungi: a source of potential antifungal compounds, *Phytochem. Rev.* 16: 883-920.
- [7] Tan, R.X., Zou, W.X. (2001) Endophytes: a rich source of functional metabolites, *Nat. Prod. Rep.* 18: 448-459.
- [8] Gunatilaka, A.A. (2006). Natural products from plant-associated microorganisms: distribution, structural diversity, bioactivity, and implications of their occurrence, *J. Nat. Prod.* 69: 509-526.
- [9] Verma, V.C., Kharwar, R.N., Strobel, G.A. (2009). Chemical and functional diversity of natural products from plant associated endophytic fungi, *Nat. Prod. Commun.* 4: 1511-1532.
- [10] Aly, A.H., Debbab, A., Kjer, J., Proksch, P. (2010). Fungal endophytes from higher plants: a prolific source of phytochemicals and other bioactive natural products, *Fungal Divers.* 41: 1-16.
- [11] Zhang, Y.Y., Han, T., Ming, Q.L., Qin, L.P. (2012). Alkaloids produced by endophytic fungi: a review, *Nat. Prod. Commun.* 7 (7): 963-968.
- [12] Nisa, H., Kamili, A.N., Nawchoo, I.A., Shafi, S., Shameem, N., Bandh, S.A. (2015). Fungal endophytes as prolific source of phytochemicals and other bioactive natural products: a review, *Microb. Pathog.* 82: 50-59.
- [13] Sudha, V., Govindaraj, R., Baskar, K., Al-Dhabi, N.A., Duraipandiyan, V. (2016). Biological properties of endophytic fungi, *Braz. Arch. Biol. Technol.* 59: 1-7.
- [14] Köberl, M., Cardinale, M., Berg, G., Ramadan, E.M., Heuer, H., Hallmann, J., Smalla, K., Adam, M. (2013). *Bacillus* and *Streptomyces* were selected as broad-spectrum antagonists against soilborne pathogens from arid areas in Egypt. *FEMS Microbiol Lett* 342:168-178..
- [15] Egamberdieva, D., Wirth, S., Li, L., Abd-Allah, E.F., Lindström, K. (2017). Microbial cooperation in the rhizosphere improves liquorice growth under salt stress. *Bioengineered* 8:433-438.
- [16] Yehia, R.S., Osman, G.H., Assaggaf, H., Salem, R., Mohamed, M.S.M. (2020). Isolation of potential antimicrobial metabolites from endophytic fungus *Cladosporium cladosporioides* from endemic plant *Zygophyllum mandavillei*. *South African Journal of Botany* 134: 296-302..
- [17] Yasuhiro, I., Shin-Suke, M., Tsuyoshi, F., Tamotsu, F. (2010). Pterocidin, a cytotoxic compound from the endophytic *Streptomyces hygroscopicus*. *J Antibiot* 59:193-195.
- [18] Hasegawa, T., Takizawa, M., Tanida, S. (1983). A rapid analysis for chemical grouping of aerobic actinomycetes. *J Gen Appl Microbiol* 29:319-322.
- [19] Kharwar, R.N., Verma, V.C., Kumar, A., Gond, S.K., Harper, J.K., Hes, W.M., Lobkovosky, E., Cong, M., Ren, Y., Strobel, G.A. (2009). Javanicin, an antibacterial naphthaquinone from an endophytic fungus of neem, *Chloridium* sp. *Curr Microbiol* 58:233-238.
- [20] Mamangkey, J., Suryanto, D., Munir, E., Lutfia, A., Hartanto, A., Huda, M.K. (2019). First Report of Plant Growth Promoting Endophytic Bacteria from Medicinal Invasive Plants (*Chromolaena odorata*). IOP Conf. Series: Earth and Environmental Science 305: 012091..

- [21] Castronovo, L.M., Vassallo. A., Mengoni, A., Miceli, E., Bogani, P., Firenzuoli, F., Fani, R., Maggini, V. (2021). Medicinal Plants and Their Bacterial Microbiota: A Review on Antimicrobial Compounds Production for Plant and Human Health. *Pathogens*, 10: 106.
- [22] Nxumalo, C.I., Ngidi, L.S., Shandu, J.S.E., Maliehe, T.S. (2020). Isolation of endophytic bacteria from the leaves of *Anredera cordifolia* CIX1 for metabolites and their biological activities. *BMC Complementary Medicine and Therapies* 20:300.
- [23] Cardoso, F.M., Campos, F.F., Santos, A.R.O., Ottoni, M.H.F., Rosa, C.A., Almeida, F.G., Graef, C.F.F. (2020). Biotechnological applications of the medicinal plant *Pseudobrickellia brasiliensis* and its isolated endophytic bacteria. 129 (4): 926-934.
- [24] Yousef, N., Mawad, A., Abeed, A. (2019). Enhancement the Cellulase Activity Induced by Endophytic Bacteria Using Calcium Nanoparticles. *Curr. Microbiol.* 76, 346-354.
- [25] Hua, L., Qi, W.Y., Hussain, S.H., Gao, K., Arfan, M. (2012). Highly oxygenated stigmastane-type steroids from the aerial parts of *Vernonia anthelmintica* Willd. *Steroids* 77:811–818.
- [26] Ito, T., Aimaiti, S., Win, N.N., Kodama, T., Morita, H. (2016). New sesquiterpene lactones, vernonilides A and B, from the seeds of *Vernonia anthelmintica* in Uyghur and their antiproliferative activities. *Bioorg Med Chem Lett* 26:3608–3611.
- [27] Yousef, N., Mawad, A., Abeed, A. (2019). Enhancement the Cellulase Activity Induced by Endophytic Bacteria Using Calcium Nanoparticles. *Curr. Microbiol.* 76, 346-354.
- [28] Fatima, S.S., Rajasekhar, M.D., Kumar, K.V., Kumar, M.T.S., Babu, K.R., Rao, C.A. (2010). Antidiabetic and antihyperlipidemic activity of ethyl acetate:isopropanol (1:1) fraction of *Vernonia anthelmintica* seeds in streptozotocin induced diabetic rats. *Food Chem Toxicol* 48:495-501.
- [29] Pasaribu, G., Budianto, E., Cahyana, H., Saepudin, E. (2020). A Review on Genus *Saurauia*: Chemical Compounds and their Biological Activity. *Pharmacogn J.* 12(3): 657-666.
- [30] Silalahi, M., Supriatna, J., Walujo, E.K.O.B. (2015). Local knowledge of medicinal plants in sub-ethnic Batak Simalungun of North Sumatra, Indonesia. *Biodiversitas* 16, 44-54.
- [31] Lentz, D.L., Clark, A.M., Hufford, C.D., Meurer-Grimes, B., Passreiter, C.M., Cordero, J., Ibrahim, O., Okunade, A.L. (1998). Antimicrobial properties of Honduran medicinal plants. *J. Ethnopharmacol.* 63, 253-263.
- [32] Yenti, R., Afrianti, R., Afriani, L. (2011). Formulasi krim ekstrak etanol daun kirinyuh (*Euphorium odoratum* L.) untuk penyembuhan luka. *Majalah Kesehatan PharmaMedika*, 3(1), 227-230.
- [33] Srivastava, R., Verma, A., Mukerjee, A., Soni, N. (2014). Phytochemical, Pharmacological and pharmacognostical profile of *Vernonia anthelmintica*: an overview. *Res Rev J Pharmacogn Phytochem* 2:22-28.
- [34] Dhiman, A.K. (2004). Banaras Ayurveda Series 2: Medicinal Plants of Uttaranchal State. Chowkhamba Sanskrit Series Office, Varanasi.
- [35] Kirtikar, K.R., Basu, B.D. (1935). *Indian Medicinal Plants, Vol. II*. Lalit Mohan Publication, Allahabad, 1347-1348.
- [36] Dogra, N.K., Kumar, S., Kumar, D. (2020). *Vernonia anthelmintica* (L.) Willd.: An ethnomedicinal, phytochemical, pharmacological and toxicological review. *J Ethnopharmacol.* 28 (256):112777.

- [37] Amir, F., Chin, K.Y. (2011). The chemical constituents and pharmacology of *Centratherrum anthelminticum*. *Int. J. PharmTech Res.* 3: 1772-1779.
- [38] Ocvirk, S., Kistler, M., Khan, S., Talukder, S.H., Hauner, H. (2013). Traditional medicinal plants used for the treatment of diabetes in rural and urban areas of Dhaka, Bangladesh- an ethnobotanical survey. *J. Ethnobiol. Ethnomed.* 9, 43.
- [39] Rustamova, N., Wubulikasimu, A., Mukhamedo, V.N., Gao, Y., Yili, D.E.A. (2020). Endophytic Bacteria Associated with Medicinal Plant *Vernonia anthelmintica*: Diversity and Characterization. *Curr Microbiol.* 77(8):1457-1465.
- [40] Emiliani, G., Mengoni, A., Maida, I., Perrin, E., Chiellini, C., Fondi, M., Gallo, E., Gori, L., Maggini, V., Vannacci, A. (2014). Linking bacterial endophytic communities to essential oils: clues from *Lavandula angustifolia* Mill. *Evid Based Complement Alternat Med*, 650905.
- [41] Janardhan, B.S., Vijayan, K. (2012). Types of endophytic bacteria associated with traditional medicinal plant *Lantana camara* Linn. *Pharmacogn J.* 4:20-23.
- [42] Mehanni, M.M., Safwat, M.S.A., Tantawy, A., Moaty, E.A. (2010). Endophytes of medicinal plants. *Acta Horti* 854:31-39.
- [43] Akinsanya, M.A., Goh, J.K., Lim, S.P., Ting, A.S.Y. (2015). Metagenomics study of endophytic bacteria in *Aloe vera* using next-generation technology. *Genom Data* 6:159-163.
- [44] Egamberdieva, D., Jabborova, D., Berg, G. (2016). Synergistic interactions between *Bradyrhizobium japonicum* and the endophyte *Stenotrophomonas rhizophila* and their effects on growth, and nodulation of soybean under salt stress. *Plant Soil* 405:35-45.
- [45] Berg, G., Grube, M., Schloter, M., Smalla, K. (2014). Unraveling the plant microbiome: looking back and future perspectives. *Front Microbiol* 5:148
- [46] Hoffman, A.M., Mayer, S.G., Strobel, G.A., Hess, W.M., Sovocool, G.W., Grange, A.H., Harper, J.K., Arif, A.M., Grant, D.M., Kelley-Swift, E.G. (2008). Purification, identification and activity of phomodione, a furandione from an endophytic *Phoma* species. *Phytochemistry*, 69: 1049-1056.
- [47] Kim, J.W., Choi, H.G., Song, J.H., Kang, K.S., Shim, S.H. (2018). Bioactive secondary metabolites from an endophytic fungus *Phoma* sp. PF2 derived from *Artemisia princeps* Pamp. *J Antibiot.* 72(3):174-177.
- [48] da Silva G.B.P.G., Silvino, K.F., Bezerra, J., Farias, T. (2017). Antimicrobial activity of *Phoma* sp. URM 7221: An endophyte from *Schinus terebinthifolius* Raddi (Anacardiaceae). *African Journal of Microbiology Research* 11(1):1-7.
- [49] Liu, Z., Chen, Y., Lian, B., Zhang, Z., Zhao, Y., Ji, Z., Lv, Y., Li, H. (2019). Comparative Study on Population Ecological Distribution and Extracellular Enzyme Activities of Endophytic Fungi in *Artemisia annua*. *Journal of Biosciences and Medicines*, 7: 94-105.
- [50] Buzugbe, H.S., Eze, P.M., Chukwunwe, Jim, C.R., Nwachukwu, C.U., Abonyi, D.O., Abba, C.C., Okoye, F.B.C., Esimone, C.O. (2018). Investigation of secondary metabolites of an endophytic fungus isolated from the leaves of *Chromolaena odorata* for possible antimicrobial and antioxidant Activities. *Pharm Chem J.* 5(6):72-77.
- [51] Gultom, E.S., Sakinah, M., Hasanah, U. (2020). Eksplorasi Senyawa Metabolit Sekunder Daun Kirinyuh (*Chromolaena odorata*) dengan GC-MS. *Jurnal Biosains* 6(1): 23-26.
- [46] Hoffman, A.M., Mayer, S.G., Strobel, G.A., Hess, W.M., Sovocool, G.W., Grange, A.H., Harper, J.K., Arif, A.M., Grant, D.M., Kelley-Swift, E.G. (2008). Purification,

identification and activity of phomodione, a furandione from an endophytic *Phoma* species. *Phytochemistry*, 69: 1049-1056.

Arman 2

ORIGINALITY REPORT

24%

SIMILARITY INDEX

13%

INTERNET SOURCES

22%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Nitty K. Dogra, Suresh Kumar, Dinesh Kumar. "Vernonia anthelmintica (L.) Willd.: An ethnomedicinal, phytochemical, pharmacological and toxicological review", *Journal of Ethnopharmacology*, 2020
Publication 2%
- 2 www.coursehero.com
Internet Source 2%
- 3 Hongying Wu, Zhenling Yan, Yecheng Deng, Zhengjun Wu, Xianglin Xu, Xiaowei Li, Xiaoxue Zhou, Haiyu Luo. "Endophytic fungi from the root tubers of medicinal plant *Stephania dielsiana* and their antimicrobial activity", *Acta Ecologica Sinica*, 2020
Publication 1%
- 4 "Phoma: Diversity, Taxonomy, Bioactivities, and Nanotechnology", Springer Science and Business Media LLC, 2022
Publication 1%
- 5 Jendri Mamangkey, Dwi Suryanto, Erman Munir, Anisa Lutfia, Adrian Hartanto, Muhammad Komarul Huda. " First Report of Plant Growth Promoting Endophytic

Bacteria from Medicinal Invasive Plants () ",
IOP Conference Series: Earth and
Environmental Science, 2019

Publication

6	tpcj.org Internet Source	1 %
7	www.frontiersin.org Internet Source	1 %
8	Soil Biology, 2015. Publication	1 %
9	WWW.mdpi.com Internet Source	1 %
10	www.academicjournals.org Internet Source	1 %
11	A Abubakar, H Yusuf, M Syukri, R Nasution, T Karma, A A Munawar, R Idroes. "Chemometric classification of geothermal and non-geothermal ethanol leaf extract of seurapoh (Chromolaena odorata Linn) using infrared spectroscopy", IOP Conference Series: Earth and Environmental Science, 2021 Publication	1 %
12	Patrícia Lunardelli Negreiros de Carvalho, Eliane de Oliveira Silva, Daniela Aparecida Chagas-Paula, Jaine Honorata Hortolan Luiz, Masaharu Ikegaki. "Importance and Implications of the Production of Phenolic	1 %

Secondary Metabolites by Endophytic Fungi: A Mini-Review", Mini-Reviews in Medicinal Chemistry, 2016

Publication

-
- | | | |
|----|---|-----|
| 13 | www.mdpi.com
Internet Source | 1 % |
| 14 | Ramy S. Yehia, Gamal H. Osman, Hamza Assaggaf, Reda Salem, Mahmoud S.M. Mohamed. "Isolation of potential antimicrobial metabolites from endophytic fungus <i>Cladosporium cladosporioides</i> from endemic plant <i>Zygophyllum mandavillei</i> ", <i>South African Journal of Botany</i> , 2020
Publication | 1 % |
| 15 | dokumen.pub
Internet Source | 1 % |
| 16 | www.scirp.org
Internet Source | 1 % |
| 17 | Gunawan Pasaribu, Emil Budianto, Antonius Herry Cahyana. " Toxicity and Total Phenolic Content of Extracts from Cultivation ", <i>IOP Conference Series: Materials Science and Engineering</i> , 2021
Publication | 1 % |
| 18 | iopscience.iop.org
Internet Source | 1 % |
| 19 | "Endophytes", Springer Science and Business Media LLC, 2021
Publication | 1 % |
-

- | | | |
|----|--|------|
| 20 | Chomcheon, P.. "Metabolites from the endophytic mitosporic Dothideomycete sp. LRUB20", <i>Phytochemistry</i> , 200901
Publication | <1 % |
| 21 | pubag.nal.usda.gov
Internet Source | <1 % |
| 22 | "Bioprospecting of Plant Biodiversity for Industrial Molecules", Wiley, 2021
Publication | <1 % |
| 23 | "Antioxidants in Plant-Microbe Interaction", Springer Science and Business Media LLC, 2021
Publication | <1 % |
| 24 | "Endophytes: Crop Productivity and Protection", Springer Science and Business Media LLC, 2017
Publication | <1 % |
| 25 | Ayushi Sharma, Bhanu Malhotra, Harsha Kharkwal, Giriraj T. Kulkarni, Nutan Kaushik. "Therapeutic agents from endophytes harbored in Asian medicinal plants", <i>Phytochemistry Reviews</i> , 2020
Publication | <1 % |
| 26 | Dilfuza Egamberdieva, Dilfuza Jabborova. "Plant microbiome: source for biologically active compounds", Elsevier BV, 2020
Publication | <1 % |
| 27 | Rosa Elvira Sánchez-Fernández, Rosalía Sánchez-Fuentes, Hiram Rangel-Sánchez, | <1 % |

Simón Hernández-Ortega et al. "Antifungal and antioomycete activities and modes of action of isobenzofuranones isolated from the endophytic fungus Hypoxylon anthochroum strain Gseg1", Pesticide Biochemistry and Physiology, 2020

Publication

28

[worldwidescience.org](https://www.worldwidescience.org)

Internet Source

<1 %

29

Haruna Yahaya Ismail, Ahmad Ali Farouq, Abdullahi Bako Rabah, Aminu Bayawa Muhammad et al. "chapter 15 Microbe-Assisted Phytoremediation of Petroleum Hydrocarbons", IGI Global, 2021

Publication

<1 %

30

R. Dey, K. K. Pal, M. Thomas, D.N. Sherathia et al. "ENDOPHYTIC MICROORGANISMS", Wiley, 2017

Publication

<1 %

31

Rubee Devi, Tanvir Kaur, Geetika Guleria, Kusam Lata Rana, Divjot Kour, Neelam Yadav, Ajar Nath Yadav, Anil Kumar Saxena. "Fungal secondary metabolites and their biotechnological applications for human health", Elsevier BV, 2020

Publication

<1 %

32

Shobhika Parmar, Qiaohong Li, Ying Wu, Xinya Li, Jinping Yan, Vijay K. Sharma, Yunlin Wei, Haiyan Li. " Endophytic fungal community of from two heavy metal-

<1 %

contaminated sites: evaluated by culture-dependent and culture-independent approaches ", Microbial Biotechnology, 2018
Publication

33 link.springer.com <1 %
Internet Source

34 www.nature.com <1 %
Internet Source

35 "Endophytes of Forest Trees", Springer Science and Business Media LLC, 2018 <1 %
Publication

36 "Microbiome in Plant Health and Disease", Springer Science and Business Media LLC, 2019 <1 %
Publication

37 Bhat Mohd Skinder, Bashir Ahmad Ganai, Abdul Hamid Wani. "Bioprospecting of endophytic fungi for antibacterial and antifungal activities", Elsevier BV, 2021 <1 %
Publication

38 Dilfuza Egamberdieva, Stephan Wirth, Undine Behrendt, Parvaiz Ahmad, Gabriele Berg. "Antimicrobial Activity of Medicinal Plants Correlates with the Proportion of Antagonistic Endophytes", Frontiers in Microbiology, 2017 <1 %
Publication

39 Joshi, S.. "QSAR study on antibacterial activity of sulphonamides and derived <1 %

Mannich bases", Bioorganic & Medicinal Chemistry Letters, 20031103

Publication

40

Komal Rizwan, Ismat Majeed, Muhammad Bilal, Tahir Rasheed, Ahmad Shakeel, Shahid Iqbal. "Phytochemistry and Diverse Pharmacology of Genus Mimosa: A Review", *Biomolecules*, 2022

Publication

<1 %

41

Marina Silalahi, Nisyawati. "An ethnobotanical study of traditional steam-bathing by the Batak people of North Sumatra, Indonesia", *Pacific Conservation Biology*, 2019

Publication

<1 %

42

Vania Delfino, Carmela Calonico, Antonella Lo Nostro, Lara Mitia Castronovo et al. "Antibacterial activity of bacteria isolated from against multidrug-resistant human pathogens ", *Future Microbiology*, 2021

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On